



Luis Angel Medina Andrade^{1*}, Arsenio Torres Delgado², Luis Fernando Perez Corona², Oscar Sanchez Moreno³, Daniel Flores Rodríguez⁴, Oscar Rodrigo Rocha Erazo⁵, Eduardo Vidrio Duarte⁶, José Daen Jonathan Cabañas Gomez⁷, Rene Cano Rodríguez⁵, Nestor Apaez Araujo⁷ and Oscar Arturo Nava Carmona⁸

¹General Surgery Department, General Zone Hospital #30, Mexican Social Security Institute, Mexico City, México

²General Surgery Department, Tulancingo General Hospital, Hidalgo, México

³Endoscopy Department, Gea Gonzalez Hospital, México City, México

⁴General Surgery Department, Veracruz High Specialty Regional Hospital ISSSTE, Veracruz, México

⁵General Surgery Department, Military School for Sanity Graduated, National Defense Secretary, México City, México

⁶General Surgery Department, Hospital Angeles Metropolitano, La Salle University, Mexico City, México

⁷General Surgery Department, Ecatepec General Hospital, México

⁸General Surgery Department, León General Hospital, México

Dates: Received: 03 December, 2016; Accepted: 12 January, 2017; Published: 19 January, 2017

***Corresponding author:** Luis Angel Medina Andrade, MD, General Surgery Department, General Zone Hospital #30, Mexican Social Security Institute, Mexico City, México, E-mail: buismedina_5@hotmail.com

Keywords: Pancreatitis; Severe pancreatitis; Cholecystitis complications; Nutritional management

<https://www.peertechz.com>

Introduction

Acute pancreatitis (AP) is a systemic immunoinflammatory response to auto-digestion of the pancreas and peri-pancreatic organs. AP is a common and life threatening disease. Annual incidence worldwide is 4.9–73 cases per 100000 people; affecting young adults, with a male: female ratio of 2.5:1. The mortality rate for pancreatitis is between 1.5% and 4.2% in large epidemiological studies, but varies according to pancreatitis severity, reaching 30% in those with infected pancreatic necrosis [1,2].

Case Report

Acute Pancreatitis, Actualization and Evidence Based Management

Abstract

Acute pancreatitis (AP) is a systemic immunoinflammatory response to auto-digestion of the pancreas and peri-pancreatic organs. It is a frequent gastrointestinal disease with an important morbidity, reaching 30% in severe cases. Different studies and reviews by international groups have developed multiple classification systems to assess the severity and address the correct management along time, identifying the better molecular markers, clinical outcome determinants and reaching conservative management as the angular piece in AP. In this review we present a compilation of the latest studies and international consensus about AP physiology, etiology, risk factors, diagnosis, severity assessment, imaging and treatment.

Pathophysiology

There are plenty of theories about the pathophysiology of AP, most of them conclude that distal ductal obstruction, irrespective of the mechanism, leads to upstream blockage of pancreatic secretion, which in turn prevents exocytosis of zymogen granules (containing digestive enzymes) from acinar cells. Consequently, the zymogen granules merge with intracellular lysosomes to form condensing or autophagic vacuoles, containing an admixture of digestive and lysosomal enzymes [3].

The activation of this enzymes, normally inactive into the pancreas, produces a proinflammatory signals cascade along the gland, with posterior release in the circulatory system and the consequently systemic inflammatory response syndrome (SIRS). The IL-1, IL-2, IL-6 and IL-8 release favors monocytes and macrophages quimiotaxis and signal amplification with TNF- α release by macrophages, with final permeability increase in different systems like vascular, gastrointestinal and the consequent bacterial translocation from gut lumen to the circulation [4].

Another theory propose bile acids as the pathogenic factor in biliary pancreatitis, when it is taken up by acinar cells from the bile acid transporters in apical and basolateral plasma membrane, leading to intracellular calcium increasing and the consequent increase in transcription of some proinflammatory mediators [5].

In the case of alcoholic pancreatitis, it increases digestive and lysosomal enzymes content in acinar cells, destabilizing

the organelles containing enzymes, increasing the probability of contact between lysosomal and digestive enzymes with the consequent high risk of intracellular activation of digestive enzymes [5,13].

Once the initial damage is established the progression and outcomes would depend on the medical management during the first 24 hours [6]. In case of limitation of the pancreatitis trigger and the correct initial management with aggressive fluid therapy, the pancreatic injury and the cytokine release could be limited, with the consequent decrease in SIRS and better outcomes. If it is not accomplished, acinar cells would develop ischemia and necrosis secondary to hypoperfusion [7]. Persistent cytokines release by necrotic tissue would increase vascular permeability favoring pulmonary effusion and respiratory distress, hypovolemia, hypotension, acute renal failure, intestinal edema and intra-abdominal hypertension [8,9].

Interstitial edema, absent peristalsis and increased gut permeability have been associated with bacterial translocation and sepsis. Intra-abdominal hypertension (IAH) and abdominal compartmental syndrome (ACS) can be developed in almost 35% of cases of severe pancreatitis, secondary to intestinal edema, ascites and retroperitoneal liquid collections, sometimes precipitated by an aggressive fluid therapy. In fact, actually IAH and ACS are considered a severity parameter [9-13].

Etiology

There are plenty of AP causes (Table 1). Alcohol and gallstones are responsible of 80% of cases. Incidence of idiopathic pancreatitis is increasing, maybe related with risk factors like obesity and metabolic syndrome, and the 57% of this idiopathic cases have been demonstrated with microlithiasis as cause after endoscopic ultrasound or magnetic resonance cholangiopancreatography [14-16]. People with gallstone disease will develop pancreatitis in 5%, and 25% of this will turn in a severe one [17]. Without definitive treatment in this cases recurrence is as high as 40% [18]. Choledocholithiasis is uncommon (20%-30%) following a mild attack of ABP [19]. Gallstones <5 mm in diameter are more likely to cause pancreatitis than larger stones. The presentation of acute pancreatitis in only 2-3% of alcoholic people suggest a genetic factor implicated [18]. In this etiology consumption of >100 g of alcohol daily and low intake of fat are significant risk factors [17].

Table 1: AP causes.

Frequent	Less Frequent
Gallstones	Autoimmune
Alcoholism	Genetic
Hypertriglyceridemia	Abdominal trauma
Post-endoscopic retrograde cholangiopancreatography	Postoperative
Idiopathic (Microlithiasis 57%)	Ischemia
Drug induced: Azathioprine, 6-Mercaptopurine, Trimethoprim-sulfamethoxazole, Pentamidine, 2,3 Dideoxyinosine, Asparaginase, Methyl-dopa.	Infections
	Hypercalcemia and hyperparathyroidism
	Posterior penetrating ulcer
	Scorpion venom
	Pancreas divisum

Risk factors

Multiple risk factors are associated with the development and severity of AP. Diabetes mellitus II have been documented to increase the risk of AP in 1.86-2.89 times [5,14,15,21].

There is a significative association between body mass index and development of biliary AP. Although there is a high prevalence of metabolic syndrome it has been demonstrated that waist circumference, body mass index, age or sex were not related with pancreatitis severity [22]. In the other hand obesity is a chronic low-grade inflammatory state characterized by high circulating levels of proinflammatory cytokines. Alternatively, obesity may intensify the immune response, which is able to exacerbate pancreatic injury and is related with a poor prognosis [13,22,23].

Current and former smoking are associated with increased risk for AP. Several experimental studies on rat models have investigated the effect of smoking showing increased inflammatory activity, focal inflammation, and decreased number of acinar structures and up-regulation of genes expressing digestive enzymes [24-26]. It has been demonstrated in studies by Sadr-Azodi, Tolstrup and Lindkvist et al., that smoking increase the relative risk of non-gallstone related acute pancreatitis 2.29, and was confirmed that duration is more important that smoking intensity⁵.

Multiple genetic factors are being studied to elucidate the pathophysiology of AP because some patients with a seemingly mild pancreatic injury (e.g., during endoscopic retrograde cholangiopancreatography [ERCP] without pancreatic duct injection) develop severe AP, whereas other subjects with extensive injury have a relatively mild course. For example, the angiotensin-converting enzyme I (rather than D) allele was significantly associated with alcohol-related AP ($p = 0.03$). The renin rs5707 G (rather than A) allele was associated with AP ($p = 0.002$), infected necrosis ($p = 0.025$) and mortality ($p = 0.046$) [27]. In preliminary studies, the authors found that the MCP-1-2518 A/G single nucleotide polymorphism predicted that the physiological response to pancreatitis would be severe and was associated with death [28]. Other mutations in genes like SPINK1, a gene that encodes for a pancreatic trypsin inhibitor, are associated with acute pancreatitis resulting from an impaired ability to counteract the effects of activated trypsin within pancreatic acinar cells.

Diagnosis and etiology assessment

The diagnosis of acute pancreatitis is based on the fulfillment of 2 of the following criteria [29]:

- Clinical upper abdominal pain
- Serum amylase or lipase >3x upper limit of normal
- Computed Tomography (CT), Magnetic Resonance Imaging (MRI), or ultrasonography diagnosis.

Once the diagnosis is established the etiology must be elucidated (Table 1), to select the correct management for better



outcomes. The main etiology is gallstones in 40% of cases. Abdominal USG is the preferred imaging study for abdominal pain associated with jaundice and for exclusion of gallstones as the cause of acute pancreatitis. Pancreas use to be visualized inadequately in 30% of cases, with about 50% of sensitivity for the detection of choledocholithiasis. Gallstones <5 mm in diameter are more likely to cause pancreatitis. An ALT>150UI/L had a positive predictive value of 95% in diagnosing acute gallstone pancreatitis [30–33].

Alcohol consumption is the responsible for about 30% of pancreatitis cases. Clinical history can elucidate the origin, like drinking more than eight alcoholic drinks/day (>100g/d) for more than 5 years. It is present in males predominantly with a male: female ratio of 2.5:1, most of them young adults [34].

Proposed mechanisms of alcohol damage include sphincter of Oddi spasm, precipitation of insoluble protein plugs that obstruct the pancreatic secretion induced by cholecystokinin [35].

Hypertriglyceridemia account for 2% of cases, and >1000mg/dL is diagnostic.

Almost 5% of Endoscopic retrograde cholangiopancreatography (ERCP) develop AP by 2 main mechanisms: traumatic intubation of the ampulla or hydrostatic pressure during contrast injection, in most of cases with a mild AP [3].

Drug induced pancreatitis is present in 2% of cases, with angiotensin-converting enzyme inhibitors, corticosteroids, diuretic and azathioprine as the principal precipitating drugs. In 0.2% of cases blunt trauma and 1% of penetrating injuries AP can be developed. Postoperative ischemia, autoimmune response, hyperparathyroidism, coxsackie, cytomegalovirus or herpes infections are less frequent causes [17].

In idiopathic acute pancreatitis 74% had biliary sludge detected by USG or had cholesterol monohydrate or calcium bilirubinate crystals detected by biliary microscopy. In cases of idiopathic pancreatitis an endoscopic USG or CPMR must be done to discard microlitiasis, present in almost 57% of cases [1]. Endoscopic ultrasound has a 90% sensitivity and 95% specificity for detecting choledocholithiasis and is somewhat more sensitive than MRCP in diagnosis of choledocholithiasis [35].

Severity assessment

Atlanta classification define three degrees of severity: mild acute pancreatitis, moderately severe acute pancreatitis, and severe acute pancreatitis, but Determinant-based classification (DBC) adds critical acute pancreatitis (Table 2) [36] and is the current classification to be employed.

This classification includes transient organ failure, persistent organ failure, and local or systemic complications. Transient organ failure is the one that is present for <48 h and persistent organ failure >48h, according to the modified Marshall score (Table 3). Local complications include peripancreatic fluid collections, pancreatic pseudocyst,

pancreatic necrotic collections and walled-off necrosis (Table 4), while systemic complications can be related to exacerbations of underlying co-morbidities exacerbated by acute pancreatitis. Mild acute pancreatitis is characterized by the absence of organ failure and the absence of local or systemic complications.

Moderately severe acute pancreatitis is characterized by the presence of transient organ failure, local or systemic complications in the absence of persistent organ failure [37]. One would expect the presence of a local complication by persistence of abdominal pain, secondary increases in serum amylase/lipase activity, organ failure, fever/chills, and so forth. Such symptoms usually prompt a cross-sectional imaging procedure to search for these complications [38].

Table 2: Classification System for severity of Acute Pancreatitis.

DBC	Mild AP	Moderate AP	Severe AP	Critical AP
Pancreatic necrosis	No	Sterile	Infected	Infected
	Both present	One or two present	Either one criterion	Both present
Organ failure	No	Transient	Persistent	Persistent

Table 3: Modified Marshall score.

System	0	1	2	3	4
Renal Creatinine(mg/dl)	<1.4	1.4-1.8	1.9-3.6	3.6-4.9	>4.9
Respiratory PaFi	>400	400-301	300-201	200-101	≤100
Cardiovascular (systolic blood pressure, mmHg) *	>90	<90 Fluid responsive	<90 Not Fluid responsive	<90 ph<7.3	<90 ph<7.2

*Without inotropic support

Organ failure is defined as a score ≥2 for one of the three scoring systems.

Multiple organ failure is defined as ≥2 systems affected.

Table 4: AP local complications.

Local complication	*CECT	Development time
Acute peripancreatic fluid collection	-Heterogeneous collection with fluid density adjacent to pancreas. -No recognizable wall encapsulating the collection. -Occurs only in interstitial edematous AP.	First 4 weeks after onset of interstitial edematous AP
Pancreatic pseudocyst	-Round or oval well circumscribed, homogeneous fluid collection. -No nonliquid component -Well-defined wall	>4 weeks after onset of interstitial edematous AP
Acute necrotic collection	-Heterogeneous nonliquid density of varying degrees -No definable encapsulating Wall -Intrapancreatic and/or extrapancreatic	Occurs in setting of acute necrotizing pancreatitis
Walled-off necrosis	-Heterogeneous liquid and nonliquid density varying degrees of loculations. -Well-defined encapsulating Wall -Intrapancreatic and/or extrapancreatic	>4 weeks after onset of necrotizing pancreatitis.

*CECT Contrast Enhanced Computed Tomography.

Severe acute pancreatitis is characterized by persistent organ failure. When SIRS is present and persistent, there is an increased risk that the pancreatitis will be complicated by persistent organ failure. Persistent organ failure may be single or multiple organ failure. Patients with persistent organ failure usually have one or more local complications. Patients who develop persistent organ failure within the first few days of the disease have an increased risk of death, with a mortality reported to be as great as 36–50%. The development of infected necrosis among patients with persistent organ failure is associated with an extremely high mortality, classified as critical acute pancreatitis [39].

Attempts to define objective criteria for assessing disease severity and prognosis were pioneered by John Ranson and Clement Imrie in the 1970s including basic laboratory data and clinical variables obtained within 48 h after hospital admission. These scoring system have found widespread application and underwent numerous modifications [40]. The Acute Physiology and Chronic Health Evaluation II (APACHE II) scoring system for critical illness may also be useful in predicting pancreatitis severity, mortality, and need for ICU admission. This was superior to both Ranson and Glasgow scores at 48 h. Although the APACHE II scoring system has gained some recognition for its performance and flexibility, the complexity of the system hinders its everyday use [18].

Abdominal hypertension (AH) and Abdominal Compartmental Syndrome (ACS) has emerged as one important parameter of severity by the relation with further complications and persistent organ failure. In a study by Ke Lu et al. the Intra-abdominal pressure (IAP), APACHE II, C-reactive protein and D-dimer was compared for the prediction of severity at 24 hours of admission. PIA and APACHE II was the more accurate to predict severe pancreatitis with a 50% higher mortality by each 1mmHg of PIA>12mmHg [9–12].

Independent markers like C reactive protein has an excellent positive predictive value for severe pancreatitis at 48 h [41–43].

In a meta-analysis of 399 patients presenting with AP, a hematocrit of >44% was predictive of the development of severe AP (along with a raised BMI and pleural effusion) [44].

Rise in blood urea nitrogen (BUN) of >5 mg/dl within 48 hours of admission was associated with the development of infected pancreatic necrosis (IPN) in 15.4% of patients, while a rise of >10 mg/dl was associated with primary IPN in 55.5% [45].

Other novel markers of severe AP include serum procalcitonin, amyloid A and cytokines such as IL-6, IL-8, IL-12 and plasma angiopoietin-2. In a multi-center study of 104 patients with predicted severe AP, a procalcitonin value of >3.5 ng/ml on two consecutive days was a more reliable marker of infected necrosis with MODS than a CRP of >430 mg/litre [46–48].

Only overweight has been related to AP severity, local complications and mortality. However, WC, BMI, sex, or age

does not correlate with disease severity [22]. Age greater than 70 has been correlated with 19% increased risk of death but is not corroborated by other studies⁴¹.

Higher morbi-mortality and interventions are needed in the AP patients with acute kidney injury, and hypertriglyceridemia is an independent risk factor for AKI. Obesity and hypertriglyceridemia increases the oxidative stress, endothelial dysfunction, inflammation and AKI [22,48].

Imaging

During initial evaluation an USG to discard gallstones as AP origin must be performed.

The gold standard for pancreas evaluation is a contrast enhanced CT scan (CECT). An early (<72h) CECT may underestimate the eventual extent of pancreatic and peripancreatic necrosis [36]. CECT is indicated only in patients with severe pancreatitis 72–96h after onset of AP, in patients with an uncertain diagnosis or when the clinical course is worsening with correct treatment, looking for local complications. It allows us to identify pancreatic lesions and define it is an edematous or necrotic pancreatitis. The local complications to be identified include infected necrosis (gas presence), walled of necrosis, pseudocyst and peripancreatic fluid collections. One week CECT for follow up is recommended to perform, with thin collimation and slice thickness (i.e. 5mm or less), 100–150 ml of non-ionic intra-venous contrast material at a rate of 3mL/s, during the pancreatic and/or portal venous phase (i.e. 50–70 seconds delay). During follow up only a portal venous phase (monophasic) is generally sufficient. For MR, the recommendation is to perform axial FS-T2 and FS-T1 scanning before and after intravenous gadolinium contrast administration [29].

Treatment

Fluid Therapy: Fluid therapy must be 5–10ml/k/h after AP onset, with lactated ringer as the identified better solution because it reduces the incidence of SIRS by 80% compared with saline resuscitation [10,29,49].

Analgesia: In a Cochrane study analyzing five randomized controlled trials the use of opioids for analgesia was associated with appropriate pain relief, decreasing the need of supplementary analgesia and no differences in complications associated with analgesics used, including nausea, vomiting, and somnolence-sedation [50].

The used opiates are preferred by their potency, but the one administered should not induce Oddi hypertension that could exacerbate pancreatitis like in the case of Morphine. Meperidine at doses of 50 to 100mg every 3 hours is safe and not associated with raise of Oddi sphincter pressure, but must be administered only for a few days because the accumulation of normeperidine metabolite can cause agitation and seizures. Somnolence or hypoventilation must be avoided by correct titration and monitoring [37].

Nutritional support: Oral feeding in predicted mild pancreatitis can be restarted once abdominal pain is decreasing.

Nutritional support is indicated 48 hours after severe AP onset. Enteral nutritional support will always be preferred. In case of not tolerating oral feeding a nasogastric or nasojejunal tube must be installed, and polymeric or elemental formulations can be used. Enteral feeding preserve physical gut barrier function, reduce microbial translocation, improve gut blood flow, preserve gut mucosal surface immunity, and maintain gut-associated lymphoid tissue mass and function [51]. These factors contribute to better outcomes and limited SIRS, less infectious complications and inclusive pain release in 25% of cases [29,49]. Cochrane studies report that enteral nutrition compared to parenteral significantly reduces mortality, multiple organ failure, systemic infections, length of hospital stay and the need for operative interventions [51].

Parenteral nutrition can be administered in acute pancreatitis as second-line therapy if nasojejunal tube feeding is not tolerated and nutritional support is required. Immunonutrients like glutamine and ω -3 fat acids added to parenteral formulas can improve prognoses in patients with acute pancreatitis. Parenteral immunonutrition significantly reduced the risk of infectious complications (RR ¼ 0.59; 95% CI, 0.39-0.88; $p= 0.05$) and mortality (RR ¼ 0.26; 95% CI, 0.11-0.59; $p= 0.001$). Length of hospital stay was also shorter in patients who received immunonutrition (MD ¼ 2.93 days; 95% CI, 4.70 to 1.15; $p=0.001$), but this results seem to be of low to very low quality [52,53].

Abdominal hypertension: Intra-abdominal hypertension (IAH) is a life-threatening sustained elevation of the intraabdominal pressure that is associated with new onset organ failure or acute worsening of existing organ failure. It is defined as >12mmHg intra-abdominal pressure. The incidence of IAH in this population is very high varying from 60 to 85% [41,54]. Abdominal compartment syndrome (ACS) is defined as a sustained intra-abdominal pressure > 20 mmHg that is associated with new onset organ failure [29].

Zhao et al., and Wu Bu et al., found that using a resuscitation protocol with only normal saline, patients had higher intra-abdominal pressure (IAP) and ACS more often, compared to patients treated with a combination of colloids and crystalloids [55,56].

The noninvasive alternative for management include: sedation, neuromuscular blockade, nasogastric decompression, and correction of a positive cumulative fluid balance [55].

Babu et al., found that percutaneous catheter drainage (PCD) resulted in sepsis reversal in almost two-thirds of the patients, and avoided open necrosectomy despite the presence of infection in the majority of the patients undergoing PCD, in about half of them [57]. If this therapeutic is not effective median laparotomy is indicated [54-58].

Biliary pancreatitis management; During admission for mild biliary pancreatitis cholecystectomy appears safe and is recommended. Interval cholecystectomy (4 weeks) after mild biliary pancreatitis is associated with a substantial risk of readmission for recurrent biliary events, especially recurrent biliary pancreatitis in 60% of cases [59].

Early laparoscopic cholecystectomy, in the first 72 hours after admission independently of symptoms is not associated with an increased risk of complications, but is related with a shorten hospital stay in patients with mild acute pancreatitis [60].

In patients with peripancreatic collections cholecystectomy should be delayed until the collections either resolve or if they persist beyond 6 weeks. If patient have undergone sphincterotomy and are fit for surgery, cholecystectomy is advised [29].

ERCP is probably indicated in biliary pancreatitis with common bile duct obstruction. Early ERCP (<24h after onset) is only indicated only in the course of biliary AP and cholangitis. ERCP should be performed within 72hours from admission when an impacted biliary stone has been demonstrated because is related with significantly reduced mortality, local and systemic complications [29,60,61].

Local complications treatment: The optimal interventional strategy for patients with suspected or confirmed infected necrotizing pancreatitis is initial image-guided percutaneous (retroperitoneal) catheter drainage or endoscopic transluminal drainage, followed, if necessary, by endoscopic or surgical necrosectomy. This must be deleted after 4 weeks with medical treatment when possible, when the necrosis has become walled-off [29].

Endoscopic trans gastric necrosectomy compares favorably with surgery [62]. Clinical trials are needed to validate the various options for intervention. Van Santvoort and colleagues compared step-up management of infected necrosis (placement of percutaneous catheters in addition to treatment with antibiotics, if necessary followed by minimally invasive necrosectomy) with open necrosectomy. This step-up approach reduced new-onset multi-organ failure by 29% [63]. Laparoscopic retroperitoneal necrosectomy is an option to avoid possible contamination of abdominal cavity and has demonstrated good outcomes.

Pseudocyst spontaneous resolution occurs in a third of patients with a pseudocyst <4cm [64]. Symptomatic pseudocysts can be successfully decompressed by endoscopic cyst gastrostomy with endoscopic ultrasound guidance [65].

Ductal disruption can result in unilateral pleural effusion, pancreatic ascites, or enlarging fluid collection, and placement of a birding stent via ERCP usually promotes duct healing if the disruption is focal [3].

Surgical management: Surgical intervention is only indicated in the course of infected necrosis, clinical deterioration after the failure of conservative management, persistent symptoms such as gastric, intestinal or biliary obstruction, pain due to the mass effect or ACS. Initial management of ACS must be medical, and if it fails, a percutaneous guided drainage must be installed. Only if this two steps fail a decompressive laparotomy must be done.

Cochrane studies report that actually low to very low quality evidence suggested that the minimally invasive step-up

approach resulted in fewer adverse events, organ failure, and total cost compared with open necrosectomy. Very low quality evidence suggested that the endoscopic minimally invasive step-up approach resulted in fewer adverse events than the video-assisted minimally invasive step-up approach but increasing the number of procedures required for treatment. In the future the TENSION trial would elucidate with a higher level of evidence which of these procedures or combination of procedures obtain the best outcomes [66].

Surgical necrosectomy, if indicated, should be done at a late stage, at least 2 weeks after the onset of pancreatitis, and only after minimally invasive methods have failed for the high morbidity associated with the procedure and poor outcomes [67]. In the course of these procedures the use of drainage is very common, and the early removal is highly recommended to reduce associated complications, length of hospital stay and total hospital cost [68].

Antibiotics

Seven evaluable studies randomized 404 patients. There was no statistically significant effect on reduction of mortality with therapy: 8.4% versus controls 14.4%, and infected pancreatic necrosis rates: 19.7% versus controls 24.4%. Non-pancreatic infection rates and the incidence of overall infections were not significantly reduced with antibiotics: 23.7% versus 36%; 37.5% versus 51.9% respectively. Operative treatment and fungal infections were not significantly different. Insufficient data were provided concerning antibiotic resistance.

With beta-lactam antibiotic prophylaxis there was less mortality (9.4% treatment vs 15% controls), and less infected pancreatic necrosis (16.8% treatment group vs 24.2% controls) but this was not statistically significant. The incidence of non-pancreatic infections was non-significantly different (21% versus 32.5%), as was the incidence of overall infections (34.4% versus 52.8%), and operative treatment rates. No significant differences were seen with quinolone plus imidazole in any of the end points measured. Imipenem on its own showed no difference in the incidence of mortality, but there was a significant reduction in the rate of pancreatic infection ($p=0.02$; RR 0.34, 95% CI 0.13 to 0.84) [69].

Conclusions

Systemic involvement is the main determinant of outcome in AP, having in mind that the pathogenesis of this disease is a dynamic process that, with the notable amount of data and recent high quality research of many groups, can be better understood, diagnosed and treated. Evolution in knowledge is supporting the systematic and conservative management as the angular piece to obtain better results, setting specific indications for each intervention in the evolution of the disease. All this progress leaves minimal invasive procedures and molecular biology as potential targets for new advances in the field.

References

1. Mariam N, Guy E, Michael RC (2015) Acute Pancreatitis: Update on management. *MJA* 202: 420-423. [Link: https://goo.gl/JZb5oZ](https://goo.gl/JZb5oZ)

2. Khanna AK, Meher S, Shashi P, Tiwari SK, Singh U, et al. (2013) Comparison of Ranson, Glasgow, MOSS, SIRS, BISAP, APACHE-II, CTSI Scores, IL-6, CRP, and Procalcitonin in Predicting Severity, Organ Failure, Pancreatic Necrosis, and Mortality in Acute Pancreatitis. *HPB Surgery* 2013: Article ID 367581. 10. [Link: https://goo.gl/j1hDiK](https://goo.gl/j1hDiK)

3. Paul L, Minoti A, Banks PA (2015) Acute Pancreatitis. 8: 1-12. [Link: https://goo.gl/t1bLdu](https://goo.gl/t1bLdu)

4. Papachristou GI, Whitcomb DC (2005) Inflammatory Markers of Disease Severity in Acute Pancreatitis. *Clin Lab Med* 25: 17-37. [Link: https://goo.gl/AQhqTt](https://goo.gl/AQhqTt)

5. Lankisch PG, Apte M, Banks PA (2015) Acute pancreatitis. *The Lancet* 386: 85-96 [Link: https://goo.gl/Wn58WD](https://goo.gl/Wn58WD)

6. O'Keefe SJ, Lee RB, Li J, et al. (2005) Trypsin secretion and turnover in patients with acute pancreatitis. *Am J Physiol Gastrointest Liver Physiol* 289: 181-187. [Link: https://goo.gl/g4Hqyr](https://goo.gl/g4Hqyr)

7. Braganza JM, Lee SH, McCloy RF, McMahon MJ (2011) Chronic Pancreatitis. *The Lancet* 377: 1184-1197. [Link: https://goo.gl/zJuD3x](https://goo.gl/zJuD3x)

8. Wang X, Yuwen Q, Xiuying P, Liang H, et al. (2013) An Evidence-Based Proposal for Predicting Organ Failure in Severe Acute Pancreatitis. *Pancreas* 8: 1255-1261. [Link: https://goo.gl/dfpXR4](https://goo.gl/dfpXR4)

9. Ke L, Tong ZH, Li WQ, Wu C, Li N, et al. (2014) Predictors of Critical Acute Pancreatitis: A Prospective Cohort Study. *Medicine* 93: 108. [Link: https://goo.gl/ggRvI2](https://goo.gl/ggRvI2)

10. Wu BU, Banks PA (2013) Clinical Management of Patients with Acute Pancreatitis. *Gastroenterology* 144: 1272-1281. [Link: https://goo.gl/6TWFbT](https://goo.gl/6TWFbT)

11. Bettina MR (2007) Outcome determinants in acute pancreatitis. *The American Journal of Surgery* 194: 39-44. [Link: https://goo.gl/9B0Y09](https://goo.gl/9B0Y09)

12. van Brunschot S, Schut AJ, Bouwense SA, Besselink MG, Bakker OJ (2014) Abdominal Compartment Syndrome in Acute Pancreatitis: A Systematic Review. *Pancreas* 43: 665-674. [Link: https://goo.gl/y1U7uK](https://goo.gl/y1U7uK)

13. Noel RA, Braun DK, Patterson RE, Bloomgren GL (2009) Increased risk of acute pancreatitis and biliary disease observed in patients with type 2 diabetes: a retrospective cohort study. *Diabetes Care* 32: 834-838. [Link: https://goo.gl/LPA6u9](https://goo.gl/LPA6u9)

14. Girman CJ, Kou TD, Cai B, Alexander CM, O'Neill EA, et al. (2010) Patients with type 2 diabetes mellitus have higher risk for acute pancreatitis compared with those without diabetes. *Diabetes Obes Metab* 12: 766-771. [Link: https://goo.gl/DDQ3Zo](https://goo.gl/DDQ3Zo)

15. Urushihara H, Taketsuna M, Liu Y, Oda E, Nakamura M, et al. (2012) Increased risk of acute pancreatitis in patients with type 2 diabetes: an observational study using a Japanese hospital database. *PLoS One* 7: e53224. [Link: https://goo.gl/73AoZu](https://goo.gl/73AoZu)

16. Marian N, Guy E, Michael RC (2015) Acute Pancreatitis: Update on Management *MJA* 202: 420-423. [Link: https://goo.gl/JZb5oZ](https://goo.gl/JZb5oZ)

17. Peter W, Ross C (2007) Acute Pancreatitis. *Surgery* 25: 49-56. [Link: https://goo.gl/1CTUJm](https://goo.gl/1CTUJm)

18. Kuo VC, Tarnasky PR (2013) Endoscopic Management of Acute Biliary Pancreatitis. *Gastrointest Endoscopy Clin N Am* 23: 749-768. [Link: https://goo.gl/gicmfv](https://goo.gl/gicmfv)

19. Neoptolemos JP (1989) The theory of persisting common bile duct stones in severe gallstone pancreatitis. *Ann R Coll Surg Engl* 71: 326-331. [Link: https://goo.gl/ou9zoN](https://goo.gl/ou9zoN)

20. Paul L, Minoti A, Banks PA (2015) Acute Pancreatitis. www.thelancet.com [Link: https://goo.gl/eSTqOh](https://goo.gl/eSTqOh)

21. Lai SW, Muo CH, Liao KF, Sung FC, Chen PC (2011) Risk of acute pancreatitis in type 2 diabetes and risk reduction on anti-diabetic drugs: a population-based cohort study in Taiwan. *Am J Gastroenterol* 106: 1697–1704. [Link: https://goo.gl/qWPuW7](https://goo.gl/qWPuW7)
22. Sawalhi S, Al-Maramhy H, Abdelrahman AI, Allah SE, Al-Jubori S (2014) Does the Presence of Obesity and/or Metabolic Syndrome Affect the Course of Acute Pancreatitis? A Prospective Study. *Pancreas* 43: 565-570. [Link: https://goo.gl/C3mIF4](https://goo.gl/C3mIF4)
23. Hong S, Qiwen B, Ying J, Wei A, Chaoyang T (2011) Body mass index and the risk and prognosis of acute pancreatitis: a meta-analysis. *Eur J Gastroenterol Hepatol* 23: 1136-1143. [Link: https://goo.gl/hR3z9K](https://goo.gl/hR3z9K)
24. Yuhara H, Ogawa M, Kawaguchi Y, Igarashi M, Mine T (2014) Smoking and Risk for Acute Pancreatitis A Systematic Review and Meta-Analysis. *Pancreas* 43: 1201-1207. [Link: https://goo.gl/hj8kkq](https://goo.gl/hj8kkq)
25. Lindkvist B, Appelros S, Manjer J, Berglund G, Borgstrom A (2008) A prospective cohort study of smoking in acute pancreatitis. *Pancreatol* 8: 63-70. [Link: https://goo.gl/73WKRR](https://goo.gl/73WKRR)
26. Tolstrup JS, Kristiansen L, Becker U, Grønbaek M (2009) Smoking and risk of acute and chronic pancreatitis among women and men: a population-based cohort study. *Arch Intern Med* 169: 603–609. [Link: https://goo.gl/bStGUj](https://goo.gl/bStGUj)
27. Skipworth JR, Nijmeijer RM, van Santvoort HC, Besselink MG, Schulz HU, et al. (2015) The Effect of Renin Angiotensin System Genetic Variants in Acute Pancreatitis. *Ann Surg* 261: 180–188. [Link: https://goo.gl/pkglwO](https://goo.gl/pkglwO)
28. Papachristou GI, Sass DA, Avula H, Janette I, Tony C, et al. (2004) The MCP-1 –2518 G Allele acts as a potent disease severity modifier in acute pancreatitis. *Pancreas* 29: 331. [Link: https://goo.gl/Mfk0hi](https://goo.gl/Mfk0hi)
29. Working Group IAP/APA Acute Pancreatitis Guidelines (2013) IAP/APA evidence-based guidelines for the management of acute pancreatitis. *Pancreatol* 13: e1-15. [Link: https://goo.gl/n2GURg](https://goo.gl/n2GURg)
30. Moolla Z, Anderson F, Thomson SR (2013) Use of amylase and alanine transaminase to predict acute gallstone pancreatitis in a population with high HIV prevalence. *World J Surg* 37: 156-161. [Link: https://goo.gl/OnZVbe](https://goo.gl/OnZVbe)
31. Liu CL, Fan ST, Lo CM, Tso WK, Wong Y, et al. (2005) Clinico-biochemical prediction of biliary cause of acute pancreatitis in the era of endoscopic ultrasonography. *Aliment Pharmacol Ther* 22: 423-431. [Link: https://goo.gl/zvJFG7](https://goo.gl/zvJFG7)
32. Tenner S, Dubner H, Steinberg W (1994) Predicting gallstone pancreatitis with laboratory parameters: a meta-analysis. *Am J Gastroenterol* 89: 1863-1866. [Link: https://goo.gl/iKl8Bg](https://goo.gl/iKl8Bg)
33. Nesvaderani Mariam, Eslick Guy, Michael RC (2015) Acute Pancreatitis: Update on management. *MJA* 202: 420-424. [Link: https://goo.gl/JZb5oZ](https://goo.gl/JZb5oZ)
34. Sekimoto M, Takada T, Kawarada Y, Hirata K, Mayumi T, et al. (2006) JPN Guidelines for the management of acute pancreatitis: epidemiology, etiology, natural history, and outcome predictors in acute pancreatitis. *J Hepatobiliary Pancreat Surg* 13: 10–24. [Link: https://goo.gl/044gsi](https://goo.gl/044gsi)
35. Cappell MS (2008) Acute Pancreatitis: Etiology, Clinical Presentation, Diagnosis, and Therapy. *Med Clin N Am* 92: 889–923. [Link: https://goo.gl/CWWGCX](https://goo.gl/CWWGCX)
36. Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, et al. (2013) Classification of acute pancreatitis—2012: revision of the Atlanta classification and definitions by international consensus 2012. *Gut* 62: 102–111. [Link: https://goo.gl/WWJbKF](https://goo.gl/WWJbKF)
37. Vege SS, Gardner TB, Chari ST, Munukuti P, Pearson RK, et al. (2009) Low mortality and high morbidity in severe acute pancreatitis without organ failure: a case for revising the Atlanta classification to include “moderately severe acute pancreatitis”. *Am J Gastroenterol* 104: 710-715. [Link: https://goo.gl/BslZV2](https://goo.gl/BslZV2)
38. Sarr MG, Banks PA, Bollen TL, Dervenis C, Gooszen HG, et al. (2013) The New Revised Classification of Acute Pancreatitis 2012. *Surg Clin N Am* 93: 549–562. [Link: https://goo.gl/FmLTfx](https://goo.gl/FmLTfx)
39. Mofidi R, Duff MD, Wigmore SJ, Madhavan KK, Garden OJ, et al. (2006) Association between early systemic inflammatory response, severity of multiorgan dysfunction and death in acute pancreatitis. *Br J Surg* 93: 738–744. [Link: https://goo.gl/ogGuW3](https://goo.gl/ogGuW3)
40. Alsfasser G, Rau BM, Klar E (2013) Scoring of human acute pancreatitis: state of the art. *Langenbecks Arch Surg* 398: 789-797. [Link: https://goo.gl/OYcvEq](https://goo.gl/OYcvEq)
41. De Waele Jan J (2014) Acute Pancreatitis. *Curr Opin Crit Care* 20:189-195. [Link: https://goo.gl/Z9ZQyL](https://goo.gl/Z9ZQyL)
42. Khanna AK, Meher S, Shashi P, Tiwari SK, Singh U, et al. (2013) Comparison of Ranson, Glasgow, MOSS, SIRS, BISAP, APACHE-II, CTSI scores, IL-6, CRP, and procalcitonin in predicting severity, organ failure, pancreatic necrosis, and mortality in acute pancreatitis. *HPB Surg* 2013: 367581. [Link: https://goo.gl/ZY50BH](https://goo.gl/ZY50BH)
43. Cardoso FS, Ricardoa LB, Oliveiraa AM, Hortaa DA, Papoila AL, et al. (2015) C-reactive protein at 24 Hours after Hospital Admission May Have Relevant Prognostic Accuracy in Acute Pancreatitis: A Retrospective Cohort Study. *GE Port J Gastroenterol* 187-189. [Link: https://goo.gl/fseOWA](https://goo.gl/fseOWA)
44. Gomatos IP, Xiaodong X, Ghaneh P, Halloran C, Raraty M, et al. (2014) Prognostic markers in acute pancreatitis. *Expert Rev Mol Diagn* 14: 333-346. [Link: https://goo.gl/SrBwHB](https://goo.gl/SrBwHB)
45. Talukdar R, Nechutova H, Clemens M, Vege SS (2013) Could rising BUN predict the future development of infected pancreatic necrosis? *Pancreatol* 13: 355-359. [Link: https://goo.gl/hmzkaz](https://goo.gl/hmzkaz)
46. Deanne B, Margaret K, Stephen P (2014) Acute pancreatitis. *Medicine* 43: 174-181. [Link: https://goo.gl/ffkxg3](https://goo.gl/ffkxg3)
47. Yang CJ, Chen J, Phillips ARJ, Windsor J, Petrov MS (2014) Predictors of severe and critical acute pancreatitis: a systematic review. *Dig Liver Dis* 46: 446-451. [Link: https://goo.gl/hmJtTi](https://goo.gl/hmJtTi)
48. Congye Wu, Lu Ke, Zhihui Tong, Baiqiang Li, Lei Zou, et al. (2014) Hypertriglyceridemia is a Risk Factor for Acute Kidney Injury in the Early Phase of Acute Pancreatitis. *Pancreas* 43: 1312-1316. [Link: https://goo.gl/DDvGw7](https://goo.gl/DDvGw7)
49. Schepers NJ, Besselink MG, van Santvoort HC, Bakker OJ, Bruno MJ, et al. (2013) Early management of acute pancreatitis. *Best Practice & Research Clinical Gastroenterology* 27: 727–743. [Link: https://goo.gl/FIhze8](https://goo.gl/FIhze8)
50. Basurto Ona X, Rigau Comas D, Urrútia G (2013) Opioids for acute pancreatitis pain (Review). *Cochrane Database of Systematic Reviews* 2013: Art. No.: CD009179. [Link: https://goo.gl/NgKSZj](https://goo.gl/NgKSZj)
51. Al-Omran M, AlBalawi ZH, Tashkandi MF, Al-Ansary LA (2010) Enteral versus parenteral nutrition for acute pancreatitis. *Cochrane Database of Systematic Reviews* 2010: Art. No.: CD002837. [Link: https://goo.gl/8zMmOZ](https://goo.gl/8zMmOZ)
52. Zhao XL, Zhu SF, Xue GJ, Li J, Liu YL, et al. (2015) Early oral refeeding based on hunger in moderate and severe acute pancreatitis: A prospective controlled, randomized clinical trial. *Nutrition* 31: 171–175. [Link: https://goo.gl/5yBmtd](https://goo.gl/5yBmtd)
53. Wu XM, Liao YW, Wang HY, Ji KQ, Li GF, et al. (2015) When to Initialize Enteral Nutrition in Patients With Severe Acute Pancreatitis? A Retrospective Review in a Single Institution Experience (2003–2013). *Pancreas* 44: 507-511. [Link: https://goo.gl/ZWd3l6](https://goo.gl/ZWd3l6)
54. Jafari T, Feizi A2, Askari G3, Fallah AA (2015) Parenteral immunonutrition in patients with acute pancreatitis: A systematic review and meta-analysis. *Clini Nutr* 34: 35-43. [Link: https://goo.gl/vnp2AL](https://goo.gl/vnp2AL)
55. Poropat G, Giljaca V, Hauser G, Štimac D (2015) Enteral nutrition formulations for acute pancreatitis. *Cochrane Database of Systematic Reviews* 2015, Issue 3. Art. No.: CD010605. [Link: https://goo.gl/am80um](https://goo.gl/am80um)



56. van Brunschot S, Schut AJ, Bouwense SA, Besselink MG, Bakker OJ, et al. (2014) Abdominal Compartment Syndrome in Acute Pancreatitis. *Pancreas* 43: 665-674. [Link: https://goo.gl/E9HH6J](https://goo.gl/E9HH6J)
57. Zhao G, Zhang JG, Wu HS, Tao J, Qin Q, et al. (2013) Effects of different resuscitation fluid on severe acute pancreatitis. *World J Gastroenterol* 19: 2044-2052. [Link: https://goo.gl/WqEn9y](https://goo.gl/WqEn9y)
58. Wu Bu (2011) Editorial: fluid resuscitation in acute pancreatitis: striking the right balance. *Am J Gastroenterol* 106: 1851-1852. [Link: https://goo.gl/3Y1z9i](https://goo.gl/3Y1z9i)
59. Babu RY, Gupta R, Kang M, Bhasin DK, Rana SS, et al. (2013) Predictors of surgery in patients with severe acute pancreatitis managed by the step-up approach. *Ann Surg* 257: 737-750. [Link: https://goo.gl/r2W3r6](https://goo.gl/r2W3r6)
60. Liu WH, Wang T, Yan HT, Chen T, Xu C, et al. (2015) Predictors of percutaneous catheter drainage (PCD) after abdominal paracentesis drainage (APD) in patients with moderately severe or severe acute pancreatitis along with fluid collections. *PLoS One* 1-21. [Link: https://goo.gl/U8523t](https://goo.gl/U8523t)
61. Cucher D, Kulvatunyou N, Green DJ, Jie T, Ong ES, et al. (2014) Gallstone Pancreatitis. *Surg Clin N Am* 94: 257-280. [Link: https://goo.gl/iWwqaY](https://goo.gl/iWwqaY)
62. Gurusamy KS, Nagendran M, Davidson BR (2013) Early versus delayed laparoscopic cholecystectomy for acute gallstone pancreatitis. *Cochrane Database of Systematic Reviews* Art. No.: CD010326. [Link: https://goo.gl/gc82jH](https://goo.gl/gc82jH)
63. De Waele Jan J (2014) Acute Pancreatitis. *Curr Opin Crit Care* 20: 189-195. [Link: https://goo.gl/Z9ZQyL](https://goo.gl/Z9ZQyL)
64. Bakker OJ, van Santvoort HC, van Brunschot S, Geskus RB, Besselink MG, et al. (2012) Endoscopic transgastric vs surgical necrosectomy for infected necrotizing pancreatitis: a randomized trial. *JAMA* 2012; 307: 1053-1061. [Link: https://goo.gl/Hn8HWL](https://goo.gl/Hn8HWL)
65. van Santvoort HC, Besselink MG, Bakker OJ, Hofker HS, Boermeester MA, et al. (2010) A step-up approach or open necrosectomy for necrotizing pancreatitis. *N Engl J Med* 362: 1491-502. [Link: https://goo.gl/kgJJJV](https://goo.gl/kgJJJV)
66. Lankisch PG, Weber-Dany B, Maisonneuve P, Lowenfels AB (2012) Pancreatic pseudocysts: prognostic factors for their development and their spontaneous resolution in the setting of acute pancreatitis. *Pancreatology* 12: 85-90. [Link: https://goo.gl/NpZb40](https://goo.gl/NpZb40)
67. Varadarajulu S, Christein JD, Tamhane A, Drelichman ER, Wilcox CM (2008) Prospective randomized trial comparing EUS and EGD for transmural drainage of pancreatic pseudocysts (with videos). *Gastrointest Endosc* 68: 1102-1111. [Link: https://goo.gl/c2V3ws](https://goo.gl/c2V3ws)
68. Tse F, Yuan Y (2012) Early routine endoscopic retrograde cholangiopancreatography strategy versus early conservative management strategy in acute gallstone pancreatitis. *Cochrane Database Syst Rev* Art. No. CD009779. [Link: https://goo.gl/lyAopE](https://goo.gl/lyAopE)
69. Italian Association for the Study of the Pancreas (AISP), Pezzilli R, Zerbi A, Campra D, Capurso G, et al. (2015) Consensus guidelines on severe acute pancreatitis. *Dig Liver Dis* 47: 532-543. [Link: https://goo.gl/SCm52u](https://goo.gl/SCm52u)
70. Cheng Y, Xia J, Lai M, Cheng N, He S, et al. (2015) Prophylactic abdominal drainage for pancreatic surgery. *Cochrane Database Syst Rev* 2015: Art. No.: CD010583. [Link: https://goo.gl/5wMxBj](https://goo.gl/5wMxBj)
71. Villatoro E, Mulla M, Larvin M (2010) Antibiotic therapy for prophylaxis against infection of pancreatic necrosis in acute pancreatitis. *Cochrane Database Syst Rev* Art No: CD002941. [Link: https://goo.gl/wJp5FX](https://goo.gl/wJp5FX)